

## PRACTICE PAPER 2

### SECTION-I

#### Straight Objective Type

This section contains 9 multiple choice question numbered 1 to 9. Each question has 4 choices (a), (b), (c) and (d), out of which ONLY ONE is correct.

#### Q1

Let a, b, c be three real numbers such that  $a < b < c$ . Let  $f(x)$  be continuous in  $[a, c]$  and differentiable in  $(a, c)$ . If  $f'(x)$  is strictly increasing in  $(a, c)$ , then

- a.  $(c - b)f(a) + (b - a)f(c) > (c - a)f(b)$
- b.  $(c - b)f(a) + (b - a)f(c) < (c - a)f(b)$
- c.  $f(a) < f(b) < f(c)$
- d. None of the above

#### Q2

The number of rational points on the ellipse  $\frac{x^2}{16} + \frac{y^2}{9} = 1$  is

- a. none
- b. two
- c. four
- d. Infinite

#### Q3

The number of non-negative continuous functions on  $[0, 1]$  satisfying  $\int_0^1 f(x) dx =$

$1, \int_0^1 xf(x) dx = a$  and  $\int_0^1 x^2 f(x) dx = a^2, (a \neq 0)$  is

- a. none
- b. two
- c. four
- d. Infinite

#### Q4

The hypotenuse of a right angled triangle passes through the point (2, 4) and the sides are along x and y-axis. The number of such triangles having area 18 units must be

- a. 1
- b. 2
- c. 3
- d. 4

#### Q5

Three numbers are drawn from the set  $\{1, 2, 3, \dots, n\}$  with replacement. The probability that their sum is  $2n$ , is

- a.  $\frac{1}{2n}$
- b.  $\frac{1}{2}$
- c.  $\frac{(n-1)(n+4)}{2n^3}$
- d.  $\frac{(n-1)(n+2)}{n^3}$

**Q6**

Among the following points there is only point from which tangents can be drawn to the ellipse  $3x^2 + 4y^2 = 12$  are perpendicular. The point must be

- a. (3, 4)
- b.  $(3, \sqrt{2})$
- c.  $(-2, \sqrt{3})$
- d. (1, 2)

**Q7**

The minimum value of  $\sin 3A + \sin 3B + \sin 3C$  in a triangle must be

- a. -1
- b. -2
- c.  $-\frac{\sqrt{3}}{2}$
- d. None of the above

**Q8**

If the perimeter of a triangle is 2 then the expression  $E = ab + bc + ac - abc - 1$

- a. is essentially positive
- b. is essentially negative
- c. may or may not be positive
- d. None of these

**Q9**

Let  $S = \sum_{r=1}^{2000} \sqrt{1 + \frac{1}{r^2} + \frac{1}{(r+1)^2}}$ . Then  $S$  must be equal to

- a.  $2000 \left(1 + \frac{1}{2001}\right)$
- b.  $2000 \left(1 + \frac{1}{1999}\right)$
- c.  $\sqrt{2000} \left(1 + \frac{1}{2001}\right)$
- d. None of these

**Section-II****Multiple Objective Type****Q10**

The values of  $a$  for which  $x^4 - 2ax^2 + x + a^2 - a = 0$  has all real roots are

- a. -1
- b. 1
- c. 2
- d. 3

**Q11**

If  $y$  is a function of  $x$  given by  $2 \log (y - 1) - \log x - \log (y - 2) = 0$ , then

- a. domain is  $[4, \infty]$
- b. domain is  $[0, \infty]$
- c. range is  $(2, \infty)$
- d. range is  $(0, \infty)$

**Q12**

If ordered pair  $(\alpha, \beta)$  where  $\alpha, \beta \in 1$  satisfy the equation  $2x^2 - 3xy - 2y^2 = 7$ , then value of  $\alpha + \beta$  can be

- a. 5
- b. 4
- c. -4
- d. 3

**Q13**

For the parabola  $y^2 = 4x$ , let  $P$  be the point of concurrency of three normals and  $S$  be the focus.

If  $\alpha_1$  be the sum of the angles made by three normals from the positive direction of x-axis and  $\alpha_2$  be the angle made by  $PS$  with the positive direction of x-axis then can be equal to

- a. 1
- b. 2
- c.  $1/2$
- d.  $3/2$

**Q14**

Let  $(\frac{p_1}{q_1}, \frac{p_2}{q_2})$  and  $(\frac{a_1}{b_1}, \frac{a_2}{b_2})$  be any two rational points on the circle  $x^2 + y^2 = 1$  where

$p_1, p_2, q_1, q_2, a_1, a_2, b_1$  and  $b_2$  are integers and H.C.F. of  $(p_1, q_1), (p_2, q_2), (a_1, b_1)$  and  $(a_2, b_2)$  is

1. Then the statements which are always correct are

- a.  $q_1 = q_2$
- b.  $p_1 = \pm 1$  or 0
- c.  $b_1 = b_2$
- d.  $a_1 = \pm 1$  or 0

**Q15**

If all the roots of the equation  $(x^2 - mx + n)(x^2 - nx + m) = 0$  are positive integers then  $m + n$  can be equal to

- a. 8
- b. 9
- c. 10
- d. 11

**Q16**

Let  $a, b, c$  be the lengths of the sides of a triangle  $ABC$  such that  $b + c \neq 1, c - b \neq 1$ . If  $\log_{b+c} a + \log_{c-b} a = 2 \log_{c+b} a \log_{c-b} a$ , then

a.  $\sin^2 A + \sin^2 B = \sin^2 C$

b.  $\tan A + \tan B = 1$

c.  $A + B = C$

d.  $\cos^2 A + \cos^2 B = 1$

**Q17**

If  $f(x) = \int_{x^n}^{\frac{dt}{\ln t}}, x > 0$  and  $n > m$ , then

a.  $f'(x) = \frac{x^{m-1}(x-1)}{\ln x}$

b.  $f(x)$  is decreasing for  $x > 1$

c.  $f(x)$  is increasing in  $(0, 1)$

d.  $f(x)$  is increasing for  $x > 1$

**Section-III****Assertion-Reason Type****Q18****Statement-1:**

If  $n$  is odd then the product  $P = (1 - i_1)(2 - i_2)(3 - i_3) \dots (n - i_n)$  where  $i_1, i_2, i_3, \dots, i_n$  are distinct integers taken from the set  $\{1, 2, 3, \dots, n\}$  is certainly even. Because

**Statement-2:**

$P$  can be zero for some choice of  $i_1, i_2, \dots, i_n$ .

a. Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1

b. Statement-1 is True, Statement-2 is True; Statement-2 is not a correct explanation for Statement-1

c. Statement-1 is True, Statement-2 is False

d. Statement-1 is True, Statement-2 is True

### Q19

#### **Statement-1:**

For any large positive integer  $n$ , the integer next to  $1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{1}{2n-1} - \frac{1}{2n}$  is 2.

#### **Statement-2:**

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots - \frac{1}{2n} = \frac{1}{n} + \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n-1}$$

### Q20

#### **Statement-1:**

If  $n$  leaves remainder 2 when divided by 3 then  $3^n - 1$  leaves remainder 8 when divided by 13.  
because

#### **Statement-2:**

$3^5 - 1 = 242$  leaves remainder 8 when divided by 13.

- a. Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- b. Statement-1 is True, Statement-2 is True; Statement-2 is not a correct explanation for Statement-1
- c. Statement-1 is True, Statement-2 is False
- d. Statement-1 is True, Statement-2 is True

### Q21

#### **Statement-2:**

$\tan^{-1} \frac{1}{5}$  is approximately equal to  $\frac{\pi}{16}$ . because

#### **Statement-1:**

If  $5 + i = \sqrt{26}(\cos \theta + i \sin \theta)$ , then  $(5 + i)^4 = 476 + 480i = 676(\cos 4\theta + i \sin 4\theta)$

- a. Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- b. Statement-1 is True, Statement-2 is True; Statement-2 is not a correct explanation for Statement-1
- c. Statement-1 is True, Statement-2 is False
- d. Statement-1 is True, Statement-2 is True

## Section-IV

### Linked Comprehension Type

#### M<sub>22-24</sub>: Paragraph for Question Nos. 22 to 24

Let  $I_{m, n} = \int_0^{\pi/2} \cos^m x \cos nx \, dx$

#### **Q22**

If  $m$  and  $n$  are non-negative integers, then

- $I_{m, n} > 0$  for all  $m$  and  $n$
- $I_{m, n} = 0$  for some  $m$  and  $n$
- $I_{m, n} < 0$  for some  $m$  and  $n$
- None of these

#### **Q23**

$\frac{I_{m, n}}{I_{m-2, n}}$  must be equal to

- $\frac{m(m-1)}{m^2+n^2}$
- $\frac{m(m-1)}{m^2-n^2}$
- $\frac{m(m-2)}{m^2-n^2}$
- None of these

#### **Q24**

$I_{n, n}$  must be equal to

- $\frac{\pi}{4}$
- $\frac{\pi}{2n}$
- $\frac{\pi}{2^{n+1}}$
- None of these

#### M<sub>25-27</sub>: Paragraph for Question Nos. 25 to 27

Let  $\alpha, \beta, \gamma$  be positive roots of the equation  $x^3 + ax^2 + bx + c = 0$ . answer the following questions

#### **Q25**

If  $c = -1/64$  then minimum value of  $\alpha + \beta + \gamma$  must be

- 1/3
- 1/4
- 1/2
- 3/4

#### **Q26**

If  $a = -1$  then maximum value of  $\alpha \beta^2 \gamma^3$  must be

- 3/2
- 1/2
- 1/432
- 1/64

**Q27**

If  $c = -1/64$  such that  $(\alpha + \beta)^3 - 27 \alpha \beta \gamma \leq 0$  then  $(a + b)$  must be equal to

- a.  $-\frac{9}{16}$   
 b.  $\frac{9}{16}$   
 c.  $-\frac{9}{32}$   
 d.  $-\frac{3}{4}$

**Section-V****Subject Type**

This section contains 3 questions. Write the answer of the questions (28-31) from the following combinations:

0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5

6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Q28**

If  $a, b, c$  are positive integers which are in increasing G.P. If  $\log_6 a + \log_6 b + \log_6 c = 6$  and  $b - a$  is a perfect cube then the numerical value of  $a + b + c$  must be equal to

**Q29**

If  $x, y, z > 0$  then minimum value of  $\frac{x^4 + y^4 + z^4}{xyz}$  is  $\sqrt{\lambda}$  the  $\lambda$  must be equal to

**Q30**

If  $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, B = \begin{bmatrix} \sin 60^\circ & \cos 60^\circ \\ -\cos 60^\circ & \sin 60^\circ \end{bmatrix}$ . Then  $(BB^T A)^{2007} = \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix}$ , the numerical quantity  $\lambda$  must be

**Q31**

If the roots of the equation  $x^2 + ax + b + 1 = 0$  are distinct positive integers, then min. value  $a^2 + b^2$  must be equal to