

## Physics

1. The slit width, when a light of wavelength  $6500\text{\AA}$  is incident on a slit, if first minima for red light is at  $300^\circ$ 
  - a)  $1 \times 10^{-6}\text{m}$
  - b)  $2.6 \times 10^{-6}\text{m}$
  - c)  $5.2 \times 10^{-6}\text{m}$
  - d)  $1.3 \times 10^{-6}\text{m}$
2. Newton's rings are observed by keeping a spherical surface of  $100\text{cm}$  radius on a plane glass plate. The wave length of light used is  $5880\text{\AA}$ . If the diameter of the 15th bright ring is  $0.59\text{cm}$ , the diameter of the 5th ring is
  - a)  $0.226\text{cm}$
  - b)  $0.556\text{cm}$
  - c)  $0.446\text{cm}$
  - d)  $0.336\text{cm}$
3. The resulting intensity after interference of two coherent waves represented by  $y_1 = a_1 \cos t$  and  $y_2 = a_2 \cos 2t$  will be
  - a)  $a_1 - a_2$
  - b)  $a_1 + a_2$
  - c)  $a_1^2 - a_2^2$
  - d)  $a_1^2 + a_2^2$
4. In a young's experiment, one of the slit is covered with a transparent sheet of thickness  $3.6 \times 10^{-3}\text{cm}$  due to which position of central fringe shifts to a position originally occupied by 30th bright fringe. The refractive index of the sheet, if  $\lambda = 6000\text{\AA}$  is
  - a) 1.5
  - b) 1.2
  - c) 1.3
  - d) 1.7
5. In young's double slit experiment with monochromatic light of wave length  $600\text{nm}$ , the distance between slits is  $10^{-3}\text{m}$ . For changing fringe width by  $3 \times 10^{-5}\text{m}$ 
  - a) the screen is moved away from the slits by  $5\text{cm}$
  - b) the screen is moved by  $3\text{cm}$  towards the slits
  - c) both (a) and (b) are correct
  - d) the screen is moved by  $3\text{cm}$  towards the slits
6. When two coherent monochromatic light beams of intensities  $I$  and  $4I$  are superimposed, what are the maximum and minimum possible intensities in the resulting beams?
  - a)  $5I$  and  $I$
  - b)  $5I$  and  $3I$
  - c)  $9I$  and  $I$
  - d)  $9I$  and  $3I$
7. In young's double slit experiment when violet light of wave length  $4358\text{\AA}$  is used, then 84 fringes are seen in the field of view, but when sodium light of certain wave length is used, then 62 fringes are seen in the field of view, the wave length of sodium light is
  - a)  $6893\text{\AA}$
  - b)  $5904\text{\AA}$
  - c)  $5523\text{\AA}$
  - d)  $6429\text{\AA}$
8. In an interference pattern the position of zeroth order maxima is  $4.8\text{mm}$  from a certain point P on the screen. The fringe width is  $0.2\text{mm}$ . The position of second maxima from point P is
  - a)  $5.1\text{mm}$
  - b)  $5\text{mm}$
  - c)  $40\text{mm}$
  - d)  $5.2\text{mm}$
9. If young's double slit experiment is performed in water,
  - a) the fringe width will decrease
  - b) the fringe width will increase
  - c) the fringe width will remain unchanged
  - d) there will be no fringe
10. The first diffraction minimum due to single slit diffraction is  $\theta$ , for a light of wave length  $5000\text{\AA}$ . If the width of the slit is  $1 \times 10^{-4}\text{cm}$ , then the value of  $\theta$  is
  - a) 300
  - b) 450
  - c) 600
  - d) 150
11. 2 non-coherent sources emit light beam of intensities  $I$  and  $4I$ . The maximum and minimum intensities in the resulting beam are
  - a)  $9I$  and  $3I$
  - b)  $9I$  and  $5I$
  - c)  $5I$  and  $I$
  - d)  $5I$  and  $3I$
12. Light propagates  $2\text{cm}$  distance in glass of refractive index 1.5 in time  $t_0$ . In the same time  $t_0$ , light propagates a distance of  $2.25\text{cm}$  in a medium. The refractive index of the medium is
  - a)  $4/3$
  - b)  $3/2$
  - c)  $8/3$
  - d) none of these
13. Two wave fronts are emitted from coherent sources of path difference between them is  $2.1\text{micron}$ . Path difference between the wave fronts at that point is  $7.692\pi$ . Wave length of light emitted by source will be
  - a)  $5386\text{\AA}$
  - b)  $5400\text{\AA}$
  - c)  $5460\text{\AA}$
  - d)  $5892\text{\AA}$
14. A spherical air bubble in water will act as
  - a) convex lens
  - b) plano convex lens
  - c) concave lens
  - d) glass plate
15. A concave lens can be used as a simple magnifier if the object lies

- a) beyond  $f$       b) within the focal length      c) between  $f$  and  $2f$       d) at  $2f$
16. For an equilateral prism the angle of minimum deviation is  $30^\circ$ . Then the refractive index of the material of the prism is  
a)  $1/2$       b)  $2$       c)  $2$       d)  $22$
17. Luminous flux is expressed in  
a) Lumen      b) Candela      c) Weber      d) Lue
18. Light travels through a glass plate of thickness  $d$ . If  $n$  is the refractive index of glass and  $c$  is the velocity of light in vacuum, the time taken by light to travel through the glass plate is  
a)  $n/cd$       b)  $nc/d$       c)  $nd/c$       d)  $ndc$
19. What is the magnification when an object is placed at  $2f$  of a convex mirror  
a)  $1/3$       b)  $2/3$       c)  $1$       d)  $3/2$
20. A tank is filled with water upto a height of  $12.5$  cm. The apparent depth of a needle at the bottom of the tank is ( $n$  of water =  $1.33$ )  
a)  $12.5$  cm      b)  $9.4$  cm      c)  $16.6$  cm      d)  $11.17$  cm
21. A man under water in a lake is viewing a boy standing on the bank of the lake. Then for him the boy appears to be  
a) shorter      b) taller      c) of the same size      d)  $16$  cm
22. A convex mirror placed at a distance of  $20$  cm from a candle forms a virtual image at the same position as that formed by a plane mirror at a distance of  $12$  cm from the candle. What is the focal length of the convex mirror?  
a)  $20$  cm      b)  $15$  cm      c)  $10$  cm      d)  $5$  cm
23. When light travels from 1 medium to another that remains unaltered is  
a) speed      b) wave length      c) frequency      d) intensity
24. The length of a telescope is  $100$  cm and magnification is  $19$ . The focal length of the objective and eye piece are  
a)  $90$  cm and  $10$  cm      b)  $85$  cm and  $1$  cm      c)  $95$  cm and  $25$  cm      d) None of the above
25. In a compound microscope the object produces a magnification  $10$  and eyepiece produces a magnification  $5$ . The overall magnification produced by the compound microscope is  
a)  $2$       b)  $5$       c)  $2$       d)  $50$
26. The colour of the sky is due to  
a) scattering of light      b) refraction of light      c) interference of light      d) reflection of light
27. An object is placed at a distance  $f/2$  from a convex lens of focal length  $f$ . The image will be at  
a)  $3f/2$ , real and inverted      b) one of the foci, virtual, double the size of the object      c)  $f/2$ , real and inverted      d)  $f$ , virtual and erect
28. Two thin convex lenses of focal length  $10$  cm and  $15$  cm are combined together, the focal length of the combination is  
a)  $25$  cm      b)  $12.5$  cm      c)  $15$  cm      d)  $6$  cm
29. The focal length of a convex lens is minimum for  
a) red      b) violet      c) blue      d) green
30. A biconvex lens of focal length  $20$  cm is cut out into two plano-convex lenses. The focal length of each part is  
a)  $10$  cm      b)  $20$  cm      c)  $30$  cm      d)  $40$  cm
31. The minimum distance between the object and its real image formed by a convex lens of focal length  $f$  is  
a)  $1.5 f$       b)  $f$       c)  $4 f$       d)  $3 f$
32. The refractive index of prism depends on  
a) angle of the prism      b) deviation produced by the prism      c) intensity of light      d) wave length of light
33. It is possible to observe total internal reflection when light travels from  
a) air to water      b) air to glass      c) water to glass      d) glass to water

34. A concave lens has focal length  $f$ . A real object placed at a distance  $f$  in front of the lens from the pole produces an image
- a) at infinity      b) at  $f$       c) at  $f/2$       d) at  $2/f$
35. The image formed by a plane mirror is
- a) real and same size as the object      b) virtual, same size as the object      c) real and magnified      d) none of these
36. The limit of resolution of the eye is one minute at a distance  $x$  from the eye. Two persons stand with a lateral separation of 3cm. To see the two persons just resolved by the naked eye,  $x$  should be about
- a) 20km      b) 15km      c) 10km      d) 30km
37. In the displacement method of measuring the focal length of a convex lens, the length of the images in the two positions of the lens between the object and the screen is 9cm and 4cm respectively. The length of the object is
- a) 6.25 cm      b) 1.5 cm      c) 6 cm      d) 36 cm
38. The refracting angle of a prism is  $A$  and the refractive index of the material of the prism  $\cot A/2$ , the angle of minimum deviation is
- a)  $180-A$       b)  $180-3A$       c)  $90-A$       d)  $180-2A$
39. A ray of light travels from vacuum into a medium of refractive index  $n$ . The angle of incidence is found to be twice the angle of refraction. The angle of incidence is
- a)  $\cos^{-1}n/2$       b)  $2\cos^{-1}n/2$       c)  $2\sin^{-1}n$       d)  $2\sin^{-1}n/2$
40. An object placed at distance 'a' from the focus of a convex lens forms its real image at a distance 'b' from the focus. The focal length of the mirror is
- a)  $ab$       b)  $ab$       c)  $a+b^2$       d)  $ab$
41. The distance between a point source of light and a screen is doubled. The intensity of light on the screen will be
- a) Four times the original value      b) half of the original value      c) two times the original value      d) one quarter the original value.
42. From the following which one is used for studying ultra violet light?
- a) prism of crown glass      b) prism of flint glass      c) prism of quartz      d) prism with combination of flint and crown glass
43. Electromagnetic waves are
- a) longitudinal waves      b) transverse waves      c) longitudinal nor transverse      d) stationary waves
44. If there are no atmosphere the average temperature on the surface of the earth would be
- a) lower      b) higher      c) same as now      d)  $00^\circ\text{C}$
45. Displacement current was first produced by
- a) Ampere      b) Henry      c) Maxwell      d) base
46. Pick out the odd one which has extremely short wave length much shorter than that of visible light and can be emitted from the nucleus of an atom.
- a) UV radiation      b) beta radiation      c)  $\gamma$  radiation      d) infra red radiation
47. The TV transmission tower in Delhi has a height of 240m. The distance upto when the broadcast can be received [taking radius of earth to be  $6.4 \times 10^6\text{m}$ ]
- a) 100 km      b) 60 km      c) 55 km      d) 50 km
48. All the members of electro magnetic spectrum have same
- a) frequency      b) velocity      c) wave length      d) wave number
49. Infra red spectrum lies between
- a) radio and micro wave region      b) visible and UV region      c) micro wave and visible region      d) UV and X-ray region
50. Choose the waves relevant to telecommunications.
- a) ultra violet      b) visible      c) infra red      d) micro waves