

**NSIT-Sample Paper 1**  
**MATHEMATICS**  
**MEMORY BASED QUESTIONS**

---

1.  $\int_0^{10} |x \times (x-1)(x-2)| dx$

- (a) 160.05                      (b) 1600.5                      (c) 16.005                      (d) none of these

**Sol: Ans [b]**

2. The value of  $\lim_{x \rightarrow 0} \frac{1 + \sin x - \cos x + \log(1-x)}{x^3}$  is

- (a) -1                      (b) 1/2                      (c) -1/2                      (d) 1

**Sol: Ans [c]**

3. The equation of tangent to the curve  $\frac{x^2}{3} - \frac{y^2}{2} = 1$  which is parallel to  $y = x$  is

- (a)  $y = x \pm 1$                       (b)  $y = x - 1/2$                       (c)  $y = x + 1/2$                       (d)  $y = 1 - x$

**Sol: Ans [a]**

4. If  $A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$  then  $A^{-1}$  is

(a)  $\begin{bmatrix} 1/a & 0 & 0 \\ 0 & 1/b & 0 \\ 0 & 0 & 1/c \end{bmatrix}$

(b)  $\begin{bmatrix} -1/a & 0 & 0 \\ 0 & -1/b & 0 \\ 0 & 0 & 1/c \end{bmatrix}$

(c)  $\begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 1/c \end{bmatrix}$

(d) none of these

**Sol: Ans [a]**

5. If  $\left| \frac{z+i}{z-i} \right| = 3$  then radius of the circle is

(a)  $\frac{2}{\sqrt{21}}$

(b)  $\frac{1}{\sqrt{21}}$

(c)  $\frac{\sqrt{21}}{2}$

(d)  $\sqrt{21}$

**Sol: Ans [c]**

6. Let  $f(x) = \cos x \cos 2x \cos 4x \cos 8x \cos 16x$ , then the value of  $f'(\pi/4)$  is

- (a)  $\sqrt{2}$  (b)  $-\sqrt{2}$  (c) 2 (d) -2

Sol: Ans [a]

7. Let  $(\sin a)x^2 + (\sin a)x + (1 - \cos a) = 0$ . The value of  $a$  for which roots of this equation are real and distinct.

- (a)  $(0, 2 \tan^{-1} 1/4)$  (b)  $(0, 2 \pi/3)$  (c)  $(0, \pi)$  (d)  $(0, 2\pi)$

Sol: Ans [a]

8. The angle of elevation of top of a tower from a point on the ground is  $30^\circ$  and it is  $60^\circ$  when it is viewed from a point located 40 m away from the initial point towards the tower. The height of the tower is

- (a)  $-20\sqrt{3}$  (b)  $\frac{\sqrt{3}}{20}$  (c)  $-\frac{\sqrt{3}}{20}$  (d)  $20\sqrt{3}$

Sol: Ans [d]

9. The summation of two unit vectors is a third unit vector, then the modulus of the difference of the unit vectors is

- (a)  $\sqrt{3}$  (b)  $1 - \sqrt{3}$  (c)  $1 + \sqrt{3}$  (d)  $-\sqrt{3}$

Sol: Ans [a]

10. A body falls freely from a point  $A$  and passes through the points  $B$  &  $C$ . Given that  $AB = 2BC$ . The ratio of the time taken by the body to cover the distances  $AB$  and  $BC$  is

- (a)  $\frac{2 + \sqrt{6}}{1}$  (b)  $\frac{2 - \sqrt{6}}{1}$  (c)  $\frac{1 - \sqrt{6}}{2}$  (d)  $\frac{1 + \sqrt{6}}{2}$

Sol: Ans [a]

11. The value of  $\sum_{r=0}^n r {}^n C_r x^r y^{n-r}$  where  $x + y = 1$  is equal to

- (a)  $1 - nx$  (b)  $1 + nx$  (c)  $-nx$  (d)  $nx$

Sol: Ans [d]

12. There is a set of  $m$  parallel lines intersecting a set of another  $n$  parallel lines in a plane. The number of parallelogrammes formed is

- (a)  ${}^{m-1}C_2 \cdot {}^{n-1}C_2$  (b)  ${}^m C_2 \cdot {}^n C_2$  (c)  ${}^{m-1}C_2 \cdot {}^n C_2$  (d)  ${}^m C_2 \cdot {}^{n-1}C_2$

Sol: Ans [b]

13. If in a trial the probability of success is twice the probability of failure. In six trials the probability of at least four successes is

- (a)  $\frac{496}{729}$  (b)  $\frac{400}{729}$  (c)  $\frac{500}{729}$  (d)  $\frac{600}{729}$

Sol: Ans [a]

14. A force vector  $m\mathbf{i} + n\mathbf{k}$  is applied to a body at a point  $P(1, 2, 3)$ . If moment of the force is perpendicular to  $3\mathbf{i} + 5\mathbf{j} + 6\mathbf{k}$  then relation between  $m$  &  $n$  is
- (a)  $n + 3m = 0$                       (b)  $n + 3m = 1$                       (c)  $n + 3m = 2$                       (d)  $n + 3m = 3$

**Sol: Ans [a]**

15. If  $S_1 = \Sigma n$ ,  $S_2 = \Sigma n^2$ ,  $S_3 = \Sigma n^3$  then the value of  $\lim_{n \rightarrow \infty} \frac{S_1 \left(1 + \frac{S_3}{8}\right)}{S_2^2}$  is equal to

- (a)  $3/32$                       (b)  $3/64$                       (c)  $9/32$                       (d)  $9/64$

**Sol: Ans [d]**

16. The greatest term in the expansion of  $(1 + 3x)^{54}$  where  $x = 1/3$  is
- (a)  $T_{28}$                       (b)  $T_{25}$                       (c)  $T_{26}$                       (d)  $T_{24}$

**Sol: Ans [a]**

17. The value of  $\lim_{x \rightarrow 0} \frac{(4^x - 1)^3}{\sin \frac{x^2}{4} \log(1 + 3x)}$  is

- (a)  $\frac{4}{3}(\ln 4)^2$                       (b)  $\frac{4}{3}(\ln 4)^3$                       (c)  $\frac{3}{2}(\ln 4)^2$                       (d)  $\frac{3}{2}(\ln 4)^3$

**Sol: Ans [b]**

18.  $\int_0^3 |x^3 + x^2 + 3x| dx$  is equal to

- (a)  $\frac{171}{2}$                       (b)  $\frac{171}{4}$                       (c)  $\frac{170}{4}$                       (d)  $\frac{170}{3}$

**Sol: Ans [b]**

19. The equation of family of a curve is  $y^2 = 4a(x + a)$  then differential equation of the family is
- (a)  $y = y' + x$                       (b)  $y = y'' + x$                       (c)  $y = 2y'x + y^2y^2$                       (d)  $y'' + y' + y^2 = 0$

**Sol: Ans [c]**

20. If A.M. of two numbers is twice of their G.M. then the ratio of greatest number to smallest number is
- (a)  $7 - 4\sqrt{3}$                       (b)  $7 + 4\sqrt{3}$                       (c) 21                      (d) 5

**Sol: Ans [b]**

21. Let  $A = \begin{bmatrix} 1 & 2 \\ -5 & 1 \end{bmatrix}$  and  $A^{-1} = xA + yI$ , then the value of  $x$  and  $y$  are

- (a)  $x = -1, y = 2$       (b)  $x = -1, y = -2$       (c)  $x = 1, y = 2$       (d)  $x = 1, y = -2$

Sol: Ans [a]

22. Let  $x^2 + y^2 - 2x - 6y + 6 = 0$  and  $x^2 + y^2 - 6x - 4y + 12 = 0$  are two circles, then equation of the circle having diameter as their common chord is

- (a)  $5x^2 + 5y^2 + 26x - 22y + 54 = 0$       (b)  $5x^2 + 5y^2 + 26x + 22y + 54 = 0$   
 (c)  $5x^2 + 5y^2 - 26x - 22y + 54 = 0$       (d)  $5x^2 + 5y^2 - 26x - 22y - 54 = 0$

Sol: Ans [c]

23. A plane  $x + y + z = -a\sqrt{3}$  touches the sphere  $2x^2 + 2y^2 + 2z^2 - 2x + 4y - 4z + 3 = 0$  then the value of  $a$  is

- (a)  $\pm \frac{1}{\sqrt{3}}$       (b)  $\frac{1}{2\sqrt{3}}$       (c)  $1 - \frac{1}{\sqrt{3}}$       (d)  $1 + \frac{1}{\sqrt{3}}$

Sol: Ans [a]

24. For what value of  $a, f(x) = -x^3 + 4ax^2 + 2x - 5$  is decreasing  $\forall x$ .

- (a) (1, 2)      (b) (3, 4)      (c) R      (d) no value of  $a$

Sol: Ans [d]

25. The common tangent of the parabolas  $y^2 = 4x$  and  $x^2 = -8y$  is

- (a)  $y = x + 2$       (b)  $y = x - 2$       (c)  $y = 2x + 3$       (d) none of these

Sol: Ans [d]

26. The solution of the differential equation  $\frac{dy}{dx} + \frac{2x}{1+x^2}y = \frac{1}{(1+x^2)^2}$  is

- (a)  $y(1-x^2) = \tan^{-1}x + c$       (b)  $y(1+x^2) = \tan^{-1}x + c$   
 (c)  $y(1+x^2)^2 = \tan^{-1}x + c$       (d)  $y(1-x^2)^2 = \tan^{-1}x + c$

Sol: Ans [b]

27. The value of  $\sum_{r=3}^{\infty} \frac{{}^r C_3 \cdot 3^r}{r!}$  is equal to

- (a)  $\frac{6e^2}{2}$       (b)  $\frac{6e^3}{2}$       (c)  $\frac{9e^2}{2}$       (d)  $\frac{9e^3}{2}$

Sol: Ans [d]

28. The value of  $\int \frac{1}{[(x-1)^3(x+2)^5]^{1/4}} dx$  is

- (a)  $\frac{4}{3} \left( \frac{x-1}{x+2} \right)^{1/4} + C$  (b)  $\frac{4}{3} \left( \frac{x+1}{x+2} \right)^{1/4} + C$  (c)  $\frac{4}{3} \left( \frac{x+1}{x-2} \right)^{1/4} + C$  (d)  $\frac{4}{3} \left( \frac{x-1}{x-2} \right)^{1/4} + C$

**Sol: Ans [a]**

29. Two vertices of a  $\Delta$  are  $(5, -3)$ ,  $(-2, 3)$  and orthocentre is  $(0, 0)$  then third vertex is

- (a)  $\left( \frac{38}{3}, -\frac{133}{9} \right)$  (b)  $\left( -\frac{38}{3}, \frac{133}{9} \right)$  (c)  $\left( -\frac{38}{3}, -\frac{133}{9} \right)$  (d) none of these

**Sol: Ans [c]**

30. Let  $\cos(2 \tan^{-1}x) = 1/2$  then the value of  $x$  is

- (a)  $\sqrt{3}$  (b)  $\frac{1}{\sqrt{3}}$  (c)  $1 - \sqrt{3}$  (d)  $1 - \frac{1}{\sqrt{3}}$

**Sol: Ans [b]**

31. If in a projectile motion range  $R$  is maximum then relation between  $H$  and  $R$  is

- (a)  $H = R/2$  (b)  $H = R/4$  (c)  $H = 2R$  (d)  $H = R/8$

**Sol: Ans [b]**

32. The foci of the conic section  $25x^2 + 16y^2 - 150x = 175$  are

- (a)  $(0, \pm 3)$  (b)  $(0, \pm 2)$  (c)  $(3, \pm 3)$  (d)  $(0, \pm 1)$

**Sol: Ans [c]**

33. A line passes through the point of intersection of the lines  $3x + y + 1 = 0$  and  $2x - y + 3 = 0$  and makes equal intercepts with axes. Then equation of the line is

- (a)  $5x + 5y - 3 = 0$  (b)  $x + 5y - 3 = 0$  (c)  $5x - y - 3 = 0$  (d)  $5x + 5y + 3 = 0$

**Sol: Ans [a]**

34. The value of  $(A \cup B \cup C) \cap (A \cap B^c \cap C^c)^c \cap C^c$  is

- (a)  $B \cap C^c$  (b)  $B^c \cap C^c$  (c)  $B \cap C$  (d)  $A \cap B \cap C$

**Sol: Ans [a]**

